



# A new case of facultative paedomorphosis in Smooth Newts, *Lissotriton vulgaris* (Caudata: Salamandridae), in Turkey

<sup>1</sup>Dilara Kızıl, <sup>1</sup>İlhan Bayryam İsmail, <sup>2</sup>Anthony Olivier, and <sup>1,\*</sup>Kerim Çiçek

<sup>1</sup>Section of Zoology, Department of Biology, Faculty of Science, Ege University, TR-35100, Bornova, Izmir, TURKEY <sup>2</sup>Tour du Valat, Centre de recherche pour la conservation des zones humides méditerranéennes, Le Sambuc, 13 200 Arles, FRANCE

**Abstract.**—A report of the occurrence of a dimorphic population of Smooth Newts in Lake Sazlı (Izmir, Western Anatolia, Turkey). During fieldwork on March 2015, 29 aquatic individuals (seven males, 22 females) were captured and of these individuals, five were paedomorphic (four males, one female). The metamorphic (83%) and paedomorphic (17%) ratio of the population is skewed to metamorphic. The mean snout-vent length (SVL) was 28.30 mm (range = 26.00–30.41), and total length (TL) was 57.66 mm (53.35–61.40) in paedomorphs. The average exterior gill length was 4.12 mm (2.64–4.71). The SVL was 39.99 mm (range = 33.44–39.93), and TL was 69.06 mm (66.19–79.17) in metamorphs. The possible reasons for the presence of facultative paedomorphosis in the population are discussed, with the dimorphic paedomorph hypothesis supported.

**Key words.** Salamander, Izmir, Lake Sazlı, Gediz Delta

**Citation:** Kızıl D, İsmail İB, Olivier A, and Çiçek K. 2016. A new case of facultative paedomorphosis in Smooth Newts, *Lissotriton vulgaris* (Caudata: Salamandridae), in Turkey. *Amphibian & Reptile Conservation* 10(1) [General Section]: 28–33 (e119).

**Copyright:** © 2016 Kızıl et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits unrestricted use for non-commercial and education purposes only, in any medium, provided the original author and the official and authorized publication sources are recognized and properly credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website <amphibian-reptile-conservation.org>.

**Received:** 11 February 2016; **Accepted:** 18 March 2016; **Published:** 22 June 2016


Facultative paedomorphosis is an environmentally-induced polymorphism that results in the coexistence of sexually mature, gilled, and fully aquatic paedomorphic individuals and transformed, metamorphic individuals in the same population (Denoël et al. 2005a). Paedomorphosis has been known for more than one century and can be either obligate or facultative in 57 urodeles species (Denoël et al. 2005a). Facultative paedomorphosis has been particularly reported in numerous European newts (e.g., Kalezić et al. 1994; Denoël 2005; Denoël et al. 2009a; Skorinov et al. 2009).

*Lissotriton vulgaris* has a high tendency to be paedomorphic (Beebee and Griffiths 2000) and there are many reports of the phenomenon throughout its distribution range (e.g., Skorinov et al. 2009; Litvinchuk et al. 1996; Litvinchuk 2001; Denoël et al. 2009b; Covaciu-Marcov et al. 2011; Stănescu et al. 2014). In Turkey, there are four records of facultative paedomorphosis in *L. vulgaris* from Thracian and Marmara and Aegean regions

of Turkey (Yılmaz 1983; Çevik et al. 1997; Çiçek and Ayaz 2011; Bozkurt et al. 2015, Fig. 1).

On March 13, 2015, during an amphibian and reptile monitoring survey in the Gediz Delta, paedomorphic Smooth Newts were observed in a population sampled at Lake Sazlı, which is located in the northeastern part of the Gediz Delta, approximately 15 km west of Menemen (38.600149°N, 26.911006°E, at about sea level, Fig. 1). The subspecies *Lissotriton vulgaris schmidtlerorum* (Raxworthy 1988) inhabits the Izmir region. The lake surface area is nearly 30 ha, and is surrounded by agricultural (corn, cotton, and wheat) areas and *Quercus* sp. dominated shrubs (Fig. 2a). Emergent aquatic vegetation of the lake includes Common Reed (*Phragmites australis*), Reed Mace (*Typha* sp.), Common Spike Rush (*Eleocharis* sp.), Tufted Sedge (*Carex* sp.), and Rushes (*Juncus* sp.) (Gediz Delta Management Plan 2007). Climatic conditions in the study area (Izmir) are mainly Mediterranean, with a mean annual temperature

**Correspondence.** E-mails: \*kerim.cicek@hotmail.com (or) kerim.cicek@ege.edu.tr (+90) 2323112409; Fax: (+90) 232388103; Postal Address: Section of Zoology, Department of Biology, Faculty of Science, Ege University, Bornova, Izmir, TR-35100, TURKEY (Corresponding author).



**Fig. 1.** The map of Gediz Delta (Izmir, Turkey). 1: Kumköy, Istanbul, 2: Küçükçekmece, Istanbul, 3: Ihsaniye, Karasu, Sakarya, 4: Lake Sülüklü, Manisa, 5: Lake İkiz, Izmir, The star shows the record (Lake Sazlı, Gediz Delta, Izmir), The dotted line shows Ramsar protected area borders.

of 18 °C (8.9–28 °C) and annual rainfall of 690 mm (20–1,430) (1950–2015, Turkish Meteorological Service 2016).

Sampling was conducted from the shore to one m depth by two persons using 45 cm radius dip nets. 29 aquatic individuals (seven males, 22 females) were captured, and of these individuals, five were found to be paedomorphic (four males, one female, Fig. 2b, c). The paedomorphs were captured 30–50 cm depth on shore of lake. The males presented sexual characteristics, including the distended cloacae, spots on the side, and a well-developed dorsal crest. The paedomorphs were easily discerned from the metamorphs by their gills. The sex of the captured individuals was recorded; snout-vent length [from the tip of the snout to the posterior of the cloacal opening], total length, and exterior gill length (mm) were measured with digital callipers to 0.01 mm precision. Three paedomorphic individuals were brought to the laboratory and kept in an aquarium (40 × 30 × 30 cm, in 25 cm natural spring water) for a brief period to be photographed. The paedomorphs were then released back to where they had been captured.

The mean snout-vent length (SVL) was 28.30 mm (range = 26.00–30.41, SE = 0.76), and total length (TL) was 57.66 mm (53.35–61.40, SE = 1.09) in paedomorphs. The average exterior gill length (GL) was 4.12 mm (2.64–4.71, SE = 0.38). The SVL was 39.99 mm (range = 33.44–39.93, SE = 0.81), and TL was 69.06 mm (66.19–79.17, SE = 1.33) in metamorphs (Table 1).

According to these data, the metamorphic (83%) and paedomorphic (17%) ratio of the population is skewed to metamorphic. This species shares its habitat with four

other amphibians: The Balkan-Anatolian Crested Newt (*Triturus ivanbureschi*), Levant Water Frog (*Pelophylax bedriagae*), Oriental Tree Frog (*Hyla orientalis*), and the Variable Toad (*Bufo variabilis*); five reptiles, the European Pond Turtle (*Emys orbicularis*), Western Caspian Turtle (*Mauremys rivulata*), Grass Snake (*Natrix natrix*), Dice Snake (*Natrix tessellata*), and East-Four-Lined Ratsnake (*Elaphe sauromates*). Fifteen fish species belonging to eight families are cited in the Gediz Delta Management Plan (2007), including the introduced exotic Western Mosquitofish (*Gambusia affinis*).

Baçoğlu et al. (1994) stated that when *L. vulgaris* larvae complete their metamorphosis, their TL can be 35 to 40 mm, adults can be 70 to 80 mm in TL in Western Anatolia. In the western Anatolian population, the adult snout-vent length ranged from 28.8 to 35.4 mm in males, and 30.5 to 36.4 mm in females, while TL ranged from 54.6 to 65.9 mm in males, and 56.1 to 66.8 mm in females (Olgun et al. 1999). The average SVL of *L. vulgaris* was 40.15 mm (34.7–43.7), 40.77 mm (35.3–44.6) in females (Colleoni et al. 2014). Stănescu et al. (2014) reported that the SVL of paedomorphs was 32.7 mm for females, 34.5 mm for males from the Danube Delta Biosphere Reserve (Romania). They also indicated metamorphs were larger than paedomorphs. Colleoni et al. (2014) reviewed sexual dimorphism in newts and found female-biased sex size dimorphism in the species. Bozkurt et al. (2015) found paedomorphic *L. v. koswigi* from (Sakarya) northwestern Turkey and they measured 31.60 mm for males and 30.06 mm for females. The authors claimed that the size of paedomorphs is larger than metamorphs. In the Montenegrin Smooth Newts paedomorphs may or may





**Fig. 2.** The general view of habitat (A) and a male paedomorphic *Lissotriton vulgaris* (B, C) from Lake Sazlı (Izmir, Turkey). The arrows show the cloaca (B) and the gills (C).

not be bigger than metamorphs in their developmental pathway (i.e., metamorphosis versus paedomorphosis), as well as their species and population (Denoël et al. 2009a).

The gill length (GL) of paedomorphs measured 2.58 to 7 mm in northwestern Turkey (Bozkurt et al., 2015), 5.88 mm (3.30–7.90) in Lake Sülüklü (western Turkey, Çiçek, and Ayaz, 2011), 6.7 mm in Romania (Covaciu-Marcov and Cicort-Lucaciu 2007), and 1.5 mm in Ukraine (Litvinchuk 2001). Our data was within the range of previous reports. GL varies among populations and might be connected to the oxygen level or other water quality parameters, but this remains to be demonstrated.

According to Semlitsch (1987), facultative paedomorphosis is controlled by environmental and

genetic factors that allow individuals to cope with habitat variation, take advantage of environmental heterogeneity in the presence of open niches, and increase their fitness. It can occur in a variety of habitats from deep oligotrophic alpine lakes to small eutrophic temporary ponds, arid areas and humid forests (Whiteman 1994; Denoël et al. 2001). In addition, paedomorphs are encountered in different regions and latitudes, independently of environmental and habitat conditions (Whiteman 1994; Denoël et al. 2001; Denoël et al. 2005a).

Three main hypotheses explain the appearance and maintenance of facultative paedomorphosis: i) the best of a bad lot, ii) the paedomorphic advantage, and iii) the dimorphic paedomorph hypothesis (Whiteman 1994). The paedomorph advantage hypothesis corresponds to the



**Table 1.** Summary statistics of paedomorphic and metamorphic *Lissotriton vulgaris* from Lake Sazlı (Izmir, Turkey).

	Paedomorphic Males			Paedomorphic Females		
	SVL (mm)	TL (mm)	GL (mm)	SVL (mm)	TL (mm)	GL (mm)
<b>Mean</b>	28.30	57.52	4.02	26.00	53.35	4.52
<b>Range</b>	26.91–30.41	55.49–61.40	2.64–4.71	—	—	—
<b>SE</b>	0.75	1.09	0.38	—	—	—
	Metamorphic Males			Metamorphic Females		
	SVL (mm)	TL (mm)	GL (mm)	SVL (mm)	TL (mm)	GL (mm)
<b>Mean</b>	37.46	68.80	—	36.95	68.48	—
<b>Range</b>	35.53–39.93	66.19–73.48	—	33.44–38.37	66.62–79.17	—
<b>SE</b>	1.08	1.54	—	0.84	0.89	—

basic model of Wilbur and Collins (1973). It predicts large, fast-growing individuals in good growing habitats to be paedomorphic, while individuals smaller than minimum size for the paedomorphs completely metamorphose to escape competition with larger paedomorphs. More unlikely, the best of a bad lot model predicts the reverse solution in poor habitats with low growth conditions. The larger larvae metamorphose, while the smallest ones keep a larval form and become reproductively mature. The latter hypothesis, the dimorphic paedomorph, suggests that the phenomenon results from the two other hypotheses according to the local conditions experienced by each individual (Whiteman 1994). The metamorphic and paedomorphic ratio of populations can exhibit variations across populations and species (Denoël et al. 2001). The fluctuations in natural populations of paedomorphic urodeles may be related to both natural and anthropogenic factors (Denoël et al. 2005b). The absence of predators and the abundance of food resources are known to favor the delay of metamorphosis and the appearance of paedomorphs (Denoël et al. 2001). At Lake Sülüklü (Çiçek and Ayaz 2011), the metamorph/paedomorph rate changes from year to year with no paedomorphs found in five samples in 2015 (K.Ç., pers. obs). The fluctuation in the population size of fish could affect the metamorph/paedomorph rate (Denoël et al. 2015). The Lake Sazlı population could favor the third hypothesis due to the observation that the size of paedomorphs is lower than metamorphs and presence of potential predators in the habitat.

Habitat is an essential key in the persistence of facultative paedomorphosis in natural populations of newts (Denoël 2005). Denoël and Ficetola (2014) compared the likelihood of multiple potential environmental determinants impacting facultative paedomorphosis. They observe that paedomorphs prefer deep ponds, with conditions favorable to aquatic breathing (high oxygen content), with no fish and surrounded by a suitable terrestrial habitat. Despite the presence of predators, Lake Sazlı has dense aquatic vegetation, abundant food sources and is surrounded by unsuitable terrestrial habitat. There is limited available shelter to hide and save terrestrial forms. Particularly, the presence of aquatic shelters has been shown to favor the coexistence between newts and fish (Winandy et

al. 2015). This might have allowed the co-occurrence of newts with fish in the studied population but more surveys are needed to explore these patterns.

Although several studies have documented the presence and the cause of facultative paedomorphosis in Europe (Denoël et al. 2005a and reference herein), data remain limited on Asian species and subspecies. Better monitoring of facultative paedomorphosis in this part of the world would help to explore hypotheses that may provide a more comprehensive understanding of this phenomenon.

**Acknowledgments.**—We would like to thank Oğuzkan Cumhuriyet and Ömer Döndüren for his help during the field trips, Dr. Lisa Ernoul, Dr. Arnaud Béchet (Tour du Valat), Charles La Via, and Dr. Mathieu Denoël for improving the English, and their valuable comments on the earlier version of this manuscript. This study is part of a project supported by Doğa Derneği (<http://www.dogadernegi.org/>) and the Tour du Valat ([www.tourduvalat.org](http://www.tourduvalat.org)). We are grateful to these organizations and the PACA Regional Council for their generous financial support.

## Literature Cited


- Baçoğlu M, Özeti N, Yılmaz, İ. 1994. Türkiye Amfibileri (The Amphibians of Turkey; Taxonomic List, Distribution, Key for Identification pp. 183–194). Ege Üniv. Fen Fak. Kitaplar Serisi No. 151, Bornova-Izmir, Turkey.
- Beebe TJC, Griffiths RA. 2000. *Amphibians and Reptiles, A Natural History of the British Herpetofauna*. Harper Collins Publishers, London, United Kingdom. 270 p.
- Bozkurt E, Olgun B, Wielstra B. 2015. First record of facultative paedomorphism in the Kosswig's newt *Lissotriton (vulgaris) kosswigi* (Freytag, 1955) (Urodela; Salamandridae), endemic to northwestern Turkey. *Turkish Journal of Zoology* 39: 1–5.
- Çevik İE, Atatür M.K, Arıkan H, Akyurtlaklı N. 1997. Occurrence of neotenic *Triturus vulgaris* (Urodela: Salamandridae) larvae in western Anatolia. *Israel Journal Zoology* 43: 301–304.
- Çiçek K, Ayaz D. 2011. New data on facultative paedo-

- morphism of the smooth newt, *Lissotriton vulgaris*, in Western Anatolia, Turkey. *Journal of Freshwater Ecology* 26: 99–103.
- Colleoni E, Denoël M, Padoa-Schioppa E, Scali S, Ficetola GF. 2014. Rensch's rule and sexual dimorphism in salamanders: Patterns and potential processes. *Journal of Zoology* 293: 143–151.
- Covaciu-Marcov SD, Cicort-Lucaciu AŞ. 2007. Notes on the presence of facultative paedomorphosis in the smooth newt *Lissotriton vulgaris* (Linnaeus, 1758) in western Romania. *North-West Journal of Zoology* 3: 53–57.
- Covaciu-Marcov SD, Sas I, Cicort-Lucaciu AŞ, Bogdan HV. 2011. *Lissotriton vulgaris* paedomorphs in southwestern Romania: Consequence of a human modified habitat? *Acta Herpetologica* 6: 15–18.
- Denoël M. 2005. Habitat partitioning in facultatively paedomorphic populations of Palmate newts *Triturus helveticus*. *Ambio* 34: 470–471.
- Denoël M, Duguet R, Džukić G, Kalezić M, Mazzotti S. 2001. Biogeography and ecology of paedomorphosis in *Triturus alpestris* (Amphibia, Caudata). *Journal of Biogeography* 28: 1,271–1,280.
- Denoël M, Džukić G, Kalezić M. 2005b. Effect of widespread fish introductions on paedomorphic newts in Europe. *Conservation Biology* 19: 162–170.
- Denoël M, Ficetola GF. 2014. Heterochrony in a complex world: Disentangling environmental processes of facultative paedomorphosis in an amphibian. *Journal of Animal Ecology* 83: 606–615.
- Denoël M, Ficetola GF, Ćirović R, Radović D, Džukić G, Kalezić ML, Vukov TD. 2009b. A multi-scale approach to facultative paedomorphosis of European newts in the Montenegrin karst: Distribution pattern, environmental variables and conservation. *Biological Conservation* 142: 509–517.
- Denoël M, Ivanović A, Džukić G, Kalezić ML. 2009a. Sexual size dimorphism in the evolutionary context of facultative paedomorphosis: Insights from European newts. *BMC Evolutionary Biology* 9: 278.
- Denoël M, Joly P, Whiteman HH. 2005a. Evolutionary ecology of facultative paedomorphosis in newts and salamanders. *Biological Reviews* 80: 663–671.
- Denoël M, Winandy L. 2015. The importance of phenotype diversity in conservation: Resilience of palmate newt morphotypes after fish removal in Larzac ponds (France). *Biological Conservation* 192: 402–408.
- Duellman WE, Trueb L. 1994. *Biology of the Amphibians*. The Johns Hopkins University Press, London, United Kingdom. 670 p.
- Gediz Delta Management Plan. 2007. Gediz Deltası Sulakalan Yönetim Planı (Gediz Delta Wetland Management Plan). TC Çevre ve Orman Bakanlığı Doğa Koruma ve Milli Parklar Genel Müdürlüğü, Sulakalanlar Şubesi Müdürlüğü, Ankara, Turkey. [In Turkish].
- Kalezić ML, Cvetković D, Djorovic A, Džukić G. 1994. Paedomorphosis and differences in life-history traits of two neighbouring crested newt (*Triturus carnifex*) populations. *Herpetological Journal* 4: 151–158.
- Litvinchuk SN. 2001. First record of paedomorphosis for the smooth newt (*Triturus vulgaris*) from Ukraine. *Russian Journal of Herpetology* 8: 77–78.
- Litvinchuk SN, Rudyk AM, Borkin LJ. 1996. Observation of paedomorphic newts (*Triturus vulgaris*) from the former Soviet Union. *Russian Journal of Herpetology* 3: 39–48.
- Olgun K, Baran İ, Tok CV. 1999. The Taxonomic Status of *Triturus vulgaris* (Linnaeus, 1758) Populations in Western Anatolia, Turkey. *Turkish Journal of Zoology* 23: 133–140.
- Semlitsch RD. 1987. Paedomorphosis in *Ambystoma talpoideum*: Effects of density, food and pond drying. *Ecology* 68: 994–1,002.
- Skorinov DV, Novikov O, Borkin LJ, Litvinchuk SN. 2009. Two New Cases of Paedomorphosis in the Caucasian Newts: *Ommatotriton ophryticus* (The First Record) and *Lissotriton vulgaris lantzi*. *Russian Journal of Herpetology* 16: 39–48.
- Stănescu F, Buhaciu E, Székely P, Székely D, Cogălniceanu D. 2011. Facultative paedomorphosis in a population of *Lissotriton vulgaris* (Amphibia: Salamandridae) from the Danube Delta Biosphere Reserve (Romania). *Turkish Journal of Zoology* 38 (1): 114–117.
- Whiteman HH. 1994. Evolution of facultative paedomorphosis in salamanders. *The Quarterly Review of Biology* 69: 205–221.
- Wilbur HM, Collins JP. 1973. Ecological aspects of amphibian metamorphosis. *Science* 182: 1,305–1,314.
- Winandy L, Darnet E, Denoël M. 2015. Amphibians forgo aquatic life in response to alien fish introduction. *Animal Behaviour* 109: 209–216.
- Yılmaz İ. 1983. Trakya kuyruklu kurbağaları üzerine morfolojik ve taksonomik bir araştırma (The morphological and taxonomical investigation on Thrace anurans). *Doğa Bilim Dergisi* 7: 119–130. [In Turkish].
- Turkish Meteorological Service. 2015. Available: <http://www.mgm.gov.tr/veridegerlendirme/il-ve-ilceler-istatistik.aspx?m = IZMIR> [Accessed: 10 February 2016].






**Dilara Kızıl** graduated with a biological sciences degree from Ege University (İzmir, Turkey) in 2015. She is studying for her Masters at Ege University, specializing in ecology. Dilara's undergraduate thesis was the herpetofauna of Gediz Delta. Currently her interests are the ecology of Turkish reptiles, focusing on turtles and tortoises. She is working with Kerim Çiçek at Ege University and Anthony Olivier of Tour Du Valat Research Center (France) on the ecology of amphibians and reptiles.



**İlhan Bayryam İsmail** is an ecologist and naturalist interested in amphibians. He graduated with a biological sciences degree from Ege University (İzmir, Turkey) in 2012, where he is earning his Master's degree in zoology with a thesis on the life history of the Levant Water Frog in Manisa, Turkey.



**Anthony Olivier** is a French biologist working at the Tour du Valat Institute. His research interests include ecology and conservation of amphibians and reptiles.



**Kerim Çiçek** is a herpetologist focused on taxonomy, biogeography, ecology, and conservation of Turkish amphibians and freshwater reptiles. He earned his B.S. in biology from Dumlupınar University in 2000 and his M.S. in zoology from Ege University in 2005. Kerim's Master thesis was on the food composition of Marsh Frog (*Rana ridibunda*) in Central Anatolia. He completed his Ph.D. thesis at Ege University in 2009, with the population dynamics of Uludağ Frog (*Rana macrocnemis*). Kerim is currently working at Ege University, Izmir, Turkey, has authored or co-authored over 70 peer-reviewed scientific publications, and is co-editor for the journal *Biharean Biologist*.